

Supplement 2

to the paper "Floating bundles and their applications"

A.V. Ershov

This paper is the supplement to the section 2 of the paper "Floating bundles and their applications" [1]. Below we study some properties of category, connected with cobordism rings of FBSP. In particular, we shall show that it is the tensor category.

In [1] the series $\mathfrak{G}(x, y) \in H[[x, y]] = \Omega_U^*(\widetilde{Gr})$, where $H = \Omega_U^*(Gr)$, was defined. Recall that it corresponds to the direct limit κ of the maps $\kappa_{k,l}: \widetilde{Gr}_{k,kl} \rightarrow \mathbb{C}P^{kl-1}$, where $\widetilde{Gr}_{k,kl}$ is the canonical FBSP over $Gr_{k,kl}$ $((k, l) = 1)$. In [1] some properties of $\mathfrak{G}(x, y)$ were studied. In particular, it was shown that

$$(\varepsilon \mathfrak{G})(x, y) = F(x, y),$$

where $\varepsilon: H \rightarrow R = \Omega_U^*(\text{pt})$ is the counit of the Hopf algebra H and $F(x, y) \in R[[x, y]]$ is the formal group of geometric cobordisms.

Let $\varphi_{k,l}$ be the map

$$\kappa_{k,l} \times \text{id}_{\widetilde{Gr}_{k,kl}}: \widetilde{Gr}_{k,kl} \rightarrow \mathbb{C}P^{kl-1} \times \widetilde{Gr}_{k,kl}.$$

The commutativity of the following diagram

$$\begin{array}{ccc} \widetilde{Gr}_{k,kl} & \xrightarrow{\varphi_{k,l}} & \mathbb{C}P^{kl-1} \times \widetilde{Gr}_{k,kl} \\ \varphi_{k,l} \downarrow & & \downarrow \text{id}_{\mathbb{C}P} \times \varphi_{k,l} \\ \mathbb{C}P^{kl-1} \times \widetilde{Gr}_{k,kl} & \xrightarrow{\text{diag}_{\mathbb{C}P} \times \text{id}_{\widetilde{Gr}}} & \mathbb{C}P^{kl-1} \times \mathbb{C}P^{kl-1} \times \widetilde{Gr}_{k,kl} \end{array} \quad (1)$$

allows us to define on the algebra $H[[x, y]]$ the structure of $R[[z]] = \Omega_U^*(\mathbb{C}P^\infty)$ -module such that z acts as the multiplication by $\mathfrak{G}(x, y)$. Let us denote this $R[[z]]$ -module by $(H[[x, y]]; \mathfrak{G}(x, y))$.

Let us consider $R[[z]] = \Omega_U^*(\mathbb{C}P^\infty)$ as a Hopf algebra. Recall that $\Delta_{R[[z]]}(z) = F(z \otimes 1, 1 \otimes z)$.

Proposition 1. $H[[x, y]]$ is the module coalgebra over $R[[z]]$, i. e. $R[[z]] \hat{\otimes}_R H[[x, y]] \rightarrow H[[x, y]]$ is the homomorphism of coalgebras.

Proof. The proof follows from the following commutative diagram $((km, ln) = 1)$:

$$\begin{array}{ccc}
 & \mathbb{C}P^{kl-1} \times \widetilde{Gr}_{k,kl} \times \mathbb{C}P^{mn-1} \times \widetilde{Gr}_{m,mn} & \\
 \swarrow & & \nwarrow \\
 \mathbb{C}P^{kl-1} \times \mathbb{C}P^{mn-1} \times \widetilde{Gr}_{k,kl} \times \widetilde{Gr}_{m,mn} & & \widetilde{Gr}_{k,kl} \times \widetilde{Gr}_{m,mn} \\
 \downarrow & \varphi_{k,l} \times \varphi_{m,n} & \downarrow \\
 \mathbb{C}P^{klmn-1} \times \widetilde{Gr}_{km,klmn} & \xleftarrow{\varphi_{km,ln}} & \widetilde{Gr}_{km,klmn} . \quad \square
 \end{array}$$

Let us consider the next commutative diagram $((km, ln) = 1)$:

$$\begin{array}{ccc}
 \widetilde{Gr_{k,kl} \times Gr_{m,mn}} & \xrightarrow{\psi_{kl,mn}} & \widetilde{Gr_{km,klmn}} \\
 \downarrow & & \downarrow \\
 Gr_{k,kl} \times Gr_{m,mn} & \xrightarrow{\phi_{kl,mn}} & Gr_{km,klmn} ,
 \end{array} \quad (2)$$

where $\widetilde{Gr_{k,kl} \times Gr_{m,mn}}$ is the FBSP over $Gr_{k,kl} \times Gr_{m,mn}$, induced by the map $\phi_{kl,mn}$ (the definition of $\phi_{kl,mn}$ was given in [1]). Clearly that the bundle $\widetilde{Gr_{k,kl} \times Gr_{m,mn}}$ (with fiber $\mathbb{C}P^{km-1} \times \mathbb{C}P^{ln-1}$) is (“external”) Segre’s product of the canonical FBSP over $Gr_{k,kl}$ and $Gr_{m,mn}$. By definition, put

$$\widetilde{Gr \times Gr} = \varinjlim_{(km,ln)=1} \widetilde{Gr_{k,kl} \times Gr_{m,mn}} ,$$

$$\psi = \varinjlim_{(km,ln)=1} \psi_{km,ln} : \widetilde{Gr \times Gr} \rightarrow \widetilde{Gr} .$$

We have the homomorphism of $R[[z]]$ -modules

$$\Psi : (H[[x, y]]; \mathfrak{G}(x, y)) \rightarrow (H \hat{\otimes}_R H[[x, y]]; (\Delta \mathfrak{G})(x, y)) ,$$

defined by the fiber map ψ (recall that Δ is the comultiplication in the Hopf algebra $H = \Omega_U^*(Gr)$). Clearly that the restriction $\Psi|_H$ coincides with Δ .

Let $\mathcal{P}^{k-1} \times_X \mathcal{Q}^{l-1}$ be a FBSP over a finite CW -complex X with fiber $\mathbb{C}P^{k-1} \times \mathbb{C}P^{l-1}$. Recall ([1]) that if k and l are sufficiently large then there exist a classifying map $f_{k,l}$ and the corresponding fiber map

$$\begin{array}{ccc} \mathcal{P}^{k-1} \times_X \mathcal{Q}^{l-1} & \rightarrow & \widetilde{Gr}_{k,kl} \\ \downarrow & & \downarrow \\ X & \xrightarrow{f_{k,l}} & Gr_{k,kl} \end{array} \quad (3)$$

which are unique up to homotopy and up to fiber homotopy respectively. Let $\mathcal{P}^{km-1} \times_X \mathcal{Q}^{ln-1}$, $(km, ln) = 1$ be Segre's product of $\mathcal{P}^{k-1} \times_X \mathcal{Q}^{l-1}$ with the trivial FBSP $X \times \mathbb{C}P^{m-1} \times \mathbb{C}P^{n-1}$. Let us pass to the direct limit

$$\mathcal{P} \times_X \mathcal{Q} = \varinjlim_i (\mathcal{P}^{km_i-1} \times_X \mathcal{Q}^{ln_i-1}),$$

where $(km_i, ln_i) = 1$, $m_i \mid m_{i+1}$, $n_i \mid n_{i+1}$, $m_i, n_i \rightarrow \infty$, as $i \rightarrow \infty$. The stable equivalence class of FBSP (see [1]) $\mathcal{P}^{k-1} \times_X \mathcal{Q}^{l-1}$ may be uniquely restored by the direct limit $\mathcal{P} \times_X \mathcal{Q}$. We have also a classifying map $f = \varinjlim_{(k,l)=1} f_{k,l}$ and the corresponding fiber map

$$\begin{array}{ccc} \mathcal{P} \times_X \mathcal{Q} & \rightarrow & \widetilde{Gr} \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Gr. \end{array} \quad (4)$$

Let us define the category $\mathfrak{FBSP}_f$ by the following way.

- (i) $\text{Ob}(\mathfrak{FBSP}_f)$ is the class of direct limits $\mathcal{P} \times_X \mathcal{Q}$ of FBSP over finite CW -complexes X (in other words, the class of stable equivalence classes of FBSP);
- (ii) $\text{Mor}_{\mathfrak{FBSP}_f}(\mathcal{P} \times_X \mathcal{Q}, \mathcal{P}' \times_Y \mathcal{Q}')$ is the set of fiber maps

$$\begin{array}{ccc} \mathcal{P} \times_X \mathcal{Q} & \rightarrow & \mathcal{P}' \times_Y \mathcal{Q}' \\ \downarrow & & \downarrow \\ X & \rightarrow & Y \end{array} \quad (5)$$

such that its restrictions to any fiber ($\cong \mathbb{C}P^\infty \times \mathbb{C}P^\infty$) are isomorphisms.

Applying the functor of unitary cobordisms Ω_U^* to an object $\mathcal{P} \times_X \mathcal{Q} \in \text{Ob}(\mathfrak{FBSPP}_f)$, we get the $R[[z]]$ -module $(A[[x, y]]; (f^*\mathfrak{G})(x, y)) \in \text{Ob}(\Omega_U^*(\mathfrak{FBSPP}_f))$, where $A = \Omega_U^*(X)$ and $f: X \rightarrow Gr$ is a classifying map for $\mathcal{P} \times_X \mathcal{Q}$. It is clear that $((\varepsilon_A \circ f^*)\mathfrak{G})(x, y) = F(x, y)$, where $\varepsilon_A: A \rightarrow R$ is the homomorphism, induced by an embedding of a point $\text{pt} \hookrightarrow X$. In other words, for any object in the category $\Omega_U^*(\mathfrak{FBSPP}_f)$ there exists the canonical morphism $(A[[x, y]]; (f^*\mathfrak{G})(x, y)) \rightarrow (R[[x, y]]; F(x, y))$.

Hence there exist the initial object $(H[[x, y]]; \mathfrak{G}(x, y))$ and the final object $(R[[x, y]]; F(x, y))$ in the category $\Omega_U^*(\mathfrak{FBSPP})$.

Let's consider a pair $(A[[x, y]]; (f^*\mathfrak{G})(x, y))$, $(B[[x, y]]; (g^*\mathfrak{G})(x, y)) \in \text{Ob}(\Omega_U^*(\mathfrak{FBSPP}_f))$, where $(B[[x, y]]; (g^*\mathfrak{G})(x, y)) = \Omega_U^*(\mathcal{P}' \times_Y \mathcal{Q}')$. Let's define their "tensor product" as the object $((A \otimes_R B)[[x, y]]; (((f^* \otimes g^*) \circ \Delta)\mathfrak{G})(x, y)) \in \text{Ob}(\Omega_U^*(\mathfrak{FBSPP}_f))$ (recall that $\Delta: H \rightarrow H \hat{\otimes}_R H$ is the comultiplication in the Hopf algebra H).

Proposition 2. *The category $\Omega_U^*(\mathfrak{FBSPP}_f)$ is the tensor category with the just defined tensor product and the unit $I = (R[[x, y]]; F(x, y))$.*

Proof. The proof is trivial. For example, the associativity axiom follows from the identity $((\Delta \otimes \text{id}_H) \circ \Delta)\mathfrak{G}(x, y) = (((\text{id}_H \otimes \Delta) \circ \Delta)\mathfrak{G})(x, y)$ which follows from the next commutative diagram $((kmt, lnu) = 1)$:

$$\begin{array}{ccc} \widetilde{Gr_{km,klmn}} \times Gr_{t,tu} & \rightarrow & \widetilde{Gr_{kmt,klmntu}} \\ \uparrow & & \uparrow \\ Gr_{k,kl} \times \widetilde{Gr_{m,mn}} \times Gr_{t,tu} & \rightarrow & Gr_{k,kl} \times \widetilde{Gr_{mt,mntu}}, \end{array} \quad (6)$$

where $Gr_{k,kl} \times \widetilde{Gr_{m,mn}} \times Gr_{t,tu}$ is external Segre's product of the canonical FBSP over $Gr_{k,kl}$, $Gr_{m,mn}$ and $Gr_{t,tu}$ (it is the bundle over $Gr_{k,kl} \times Gr_{m,mn} \times Gr_{t,tu}$ with fiber $\mathbb{C}P^{kmt-1} \times \mathbb{C}P^{lnu-1}$). $\square$

Note that there exist the canonical homomorphisms p_1, p_2 :

$$\begin{array}{ccc} & ((A \otimes_R B)[[x, y]]; (((f^* \otimes g^*) \circ \Delta)\mathfrak{G})(x, y)) & \\ p_1 \swarrow & & \searrow p_2 \\ (A[[x, y]]; (f^*\mathfrak{G})(x, y)) & & (B[[x, y]]; (g^*\mathfrak{G})(x, y)), \end{array}$$

such that $p_1|_{A \otimes_R B} = \text{id}_A \otimes \varepsilon_B$, $p_2|_{A \otimes_R B} = \varepsilon_A \otimes \text{id}_B$.

References

- [1] A. V. ERSHOV Floating bundles and their applications.—
arXiv:math.AT/0102054